

IGRAs: CAN THEY REPLACE THE TST?
NATIONAL WEB-BASED SEMINAR – JANUARY 28, 2009
SAN FRANCISCO, CALIFORNIA

REFERENCES:

GENERAL

WHO Report 2007, Global tuberculosis control - surveillance, planning, financing.
WHO/HTM/TB/2007.376.

Speaker #1

Introduction to IGRAs.....Neil Schluger

Bennett DE, Courval JM, Onorato I, Agerton T, Gibson JD, Lambert L, McQuillan GM, Lewis B, Navin TR, Castro KG, Prevalence of Tuberculosis Infection in the United States Population: The National Health and Nutrition Examination Survey, 1999–2000. *Am. J. Respir. Crit. Care Med.* 177: 348-355, 2008.

Mancuso JD, Tobler SK, Keep LW, Pseudoepidemics of Tuberculin Skin Test Conversions in the U.S. Army after Recent Deployments . *Am. J. Respir. Crit. Care Med.* 177: 1285-1289, 2008.

Menzies D, Pai M, Comstock G. Meta-analysis: new tests for the diagnosis of latent tuberculosis infection: areas of uncertainty and recommendations for research. *Ann Intern Med.* 146(5):340-54, 2007.

Pai M, Menzies D. The new IGRA and the old TST: making good use of disagreement. *Am J Respir Crit Care Med.* 175(6):529-31, 2007.

Schluger, NW, Assessing Tuberculosis Transmission and Virulence: The Vanishing Tuberculin Skin Test (editorial). *Am J Respir Crit Care Med.* 173: 942-943, 2006.

Mazurek GH, Jereb J, LoBue P, Iademarco MF, Metchock B, Vernon A. Guidelines for Using the QuantiFERON®-TB Gold Test for Detecting Mycobacterium tuberculosis Infection, United States. *MMWR* December 16, 2005 / 54(RR15);49-55, 2005.

Speaker #2

IGRA Use in Special Situations: Review of Currently Available Data.....Charles Daley

Lighter J, Rigaud M, Eduardo R, et al: Latent tuberculosis diagnosis in children by using the QuantiFERON-TB Gold In-Tube test. *Pediatrics* 123(1):30-37, 2009.

Nicol MP, Davies MA, Wood K, et al: Comparison of T-SPOT.TB assay and tuberculin skin test for the evaluation of young children at high risk for tuberculosis in a community setting. *Pediatrics* 123(1):38-43, 2009.

Bakir M, Millington KA, Soysal A, et al: Prognostic value of a T-cell-based, interferon-gamma biomarker in children with tuberculosis contact. *Ann Intern Med* 149(11):777-787, 2008.

Bocchino M, Matarese A, Bellofiore B, et al: Performance of two commercial blood IFN-gamma release assays for the detection of Mycobacterium tuberculosis infection in patient candidates for anti-TNF-alpha treatment. *Eur J Clin Microbiol Infect Dis* 27(10):907-913, 2008.

Chen DY, Shen GH, Hsieh TY, et al: Effectiveness of the combination of a whole-blood interferon-gamma assay and the tuberculin skin test in detecting latent tuberculosis infection in rheumatoid arthritis patients receiving adalimumab therapy. *Arthritis Rheum* 59(6):800-806, 2008.

Chun JK, Kim CK, Kim HS, et al: The role of a whole blood interferon-gamma assay for the detection of latent tuberculosis infection in Bacille Calmette-Guerin vaccinated children. *Diagn Microbiol Infect Dis* 62(4):389-394, 2008.

Connell TG, Ritz N, Paxton GA, et al: A three-way comparison of tuberculin skin testing, QuantiFERON-TB gold and T-SPOT.TB in children. *PLoS ONE* 3(7):e2624, 2008.

Diel R, Loddenkemper R, Meywald-Walter K, et al: Predictive value of a whole blood IFN-gamma assay for the development of active tuberculosis disease after recent infection with Mycobacterium tuberculosis. *Am J Respir Crit Care Med* 177(10):1164-1170, 2008.

Hill PC, Jackson-Sillah DJ, Fox A, et al: Incidence of tuberculosis and the predictive value of ELISPOT and Mantoux tests in Gambian case contacts. *PLoS ONE* 3(1):e1379, 2008.

Lindemann M, Dioury Y, Beckebaum S, et al: Diagnosis of tuberculosis infection in patients awaiting liver transplantation. *Hum Immunol*, 2008.

Pai M, Zwerling A, Menzies D: Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection: an update. *Ann Intern Med* 149(3):177-184, 2008.

Ponce de Leon D, Acevedo-Vasquez E, Alvizuri S, et al: Comparison of an interferon-gamma assay with tuberculin skin testing for detection of tuberculosis (TB) infection in patients with rheumatoid arthritis in a TB-endemic population. *J Rheumatol* 35(5):776-781, 2008.

Raby E, Moyo M, Devendra A, et al: The effects of HIV on the sensitivity of a whole blood IFN-gamma release assay in Zambian adults with active tuberculosis. *PLoS ONE* 3(6):e2489, 2008.

Stephan C, Wolf T, Goetsch U, et al: Comparing QuantiFERON-tuberculosis gold, T-SPOT tuberculosis and tuberculin skin test in HIV-infected individuals from a low prevalence tuberculosis country. *Aids* 22(18):2471-2479, 2008.

Vassilopoulos D, Stamoulis N, Hadziyannis E, et al: Usefulness of enzyme-linked immunospot assay (Elispot) compared to tuberculin skin testing for latent tuberculosis screening in rheumatic patients scheduled for anti-tumor necrosis factor treatment. *J Rheumatol* 35(7):1271-1276, 2008.

Winthrop KL, Nyendak M, Calvet H, et al: Interferon-gamma release assays for diagnosing mycobacterium tuberculosis infection in renal dialysis patients. *Clin J Am Soc Nephrol* 3(5):1357-1363, 2008.

Arend SM, Thijsen SF, Leyten EM, et al: Comparison of two interferon-gamma assays and tuberculin skin test for tracing tuberculosis contacts. *Am J Respir Crit Care Med* 175(6):618-627, 2007.

Detjen AK, Keil T, Roll S, et al: Interferon-gamma release assays improve the diagnosis of tuberculosis and nontuberculous mycobacterial disease in children in a country with a low incidence of tuberculosis. *Clin Infect Dis* 45(3):322-328, 2007.

Franken WP, Koster BF, Bossink AW, et al: Follow-up study of tuberculosis-exposed supermarket customers with negative tuberculin skin test results in association with positive gamma interferon release assay results. *Clin Vaccine Immunol* 14(9):1239-1241, 2007.

Jones S, de Gijzel D, Wallach FR, et al: Utility of QuantiFERON-TB Gold in-tube testing for latent TB infection in HIV-infected individuals. *Int J Tuberc Lung Dis* 11(11):1190-1195, 2007.

Kobashi Y, Mouri K, Obase Y, et al: Clinical evaluation of QuantiFERON TB-2G test for immunocompromised patients. *Eur Respir J* 30(5):945-950, 2007.

Luetkemeyer AF, Charlebois ED, Flores LL, et al: Comparison of an interferon-gamma release assay with tuberculin skin testing in HIV-infected individuals. *Am J Respir Crit Care Med* 175(7):737-742, 2007.

Menzies D, Pai M, Comstock G: Meta-analysis: new tests for the diagnosis of latent tuberculosis infection: areas of uncertainty and recommendations for research. *Ann Intern Med* 146(5):340-354, 2007.

Passalent L, Khan K, Richardson R, et al: Detecting latent tuberculosis infection in hemodialysis patients: a head-to-head comparison of the T-SPOT.TB test, tuberculin skin test, and an expert physician panel. *Clin J Am Soc Nephrol* 2(1):68-73, 2007.

Pratt A, Nicholl K, Kay L: Use of the QuantiFERON TB Gold test as part of a screening programme in patients with RA under consideration for treatment with anti-TNF-alpha agents: the Newcastle (UK) experience. *Rheumatology (Oxford)* 46(6):1035-1036, 2007.

Rangaka MX, Wilkinson KA, Seldon R, et al: Effect of HIV-1 infection on T-Cell-based and skin test detection of tuberculosis infection. *Am J Respir Crit Care Med* 175(5):514-520, 2007.

Vincenti D, Carrara S, Butera O, et al: Response to region of difference 1 (RD1) epitopes in human immunodeficiency virus (HIV)-infected individuals enrolled with suspected active tuberculosis: a pilot study. *Clin Exp Immunol* 150(1):91-98, 2007.

Brock I, Ruhwald M, Lundgren B, et al: Latent tuberculosis in HIV positive, diagnosed by the M. tuberculosis specific interferon-gamma test. *Respir Res* 7:56, 2006.

Diel R, Nienhaus A, Lange C, et al: Tuberculosis contact investigation with a new, specific blood test in a low-incidence population containing a high proportion of BCG-vaccinated persons. *Respir Res* 7:77, 2006.

Speaker #3

Serial Testing and Use of Quantitative Results.....Randall Reves

Pai M, Joshi R, Dogra S, Zwerling AA, Gajalakshmi D, Goswami K, Reddy MV, Kalantri A, Hill PC, Menzies D, Hopewell PC. T-cell assay conversions and reversions among household contacts of tuberculosis patients in rural India. *Int J Tuberc Lung Dis.* 13(1):84-92, 2009.

Davidow AL, Affouf M. Making sense of agreement among interferon-gamma release assays and tuberculosis skin testing. *Int J Tuberc Lung Dis.* 12(2):152-9, 2008.

Hill PC, Jackson-Sillah DJ, Fox A, Brookes RH, de Jong BC, Lugos MD, Adetifa IM, Donkor SA, Aiken AM, Howie SR, Corrah T, McAdam KP, Adegbola RA. Incidence of tuberculosis and the predictive value of ELISPOT and Mantoux tests in Gambian case contacts. *PLoS ONE.* 3(1):e1379, 2008.

Perry S, Sanchez L, Yang S, Agarwal Z, Hurst P, Parsonnet J. Reproducibility of QuantiFERON-TB gold in-tube assay. *Clin Vaccine Immunol.* 15(3):425-32, 2008.

Veerapathran A, Joshi R, Goswami K, Dogra S, Moodie EE, Reddy MV, Kalantri S, Schwartzman K, Behr MA, Menzies D, Pai M. T-cell assays for tuberculosis infection: deriving cut-offs for conversions using reproducibility data. *PLoS ONE.* 3(3):e1850, 2008.

Hill PC, Brookes RH, Fox A, Jackson-Sillah D, Jeffries DJ, Lugos MD, Donkor SA, Adetifa IM, de Jong BC, Aiken AM, Adegbola RA, McAdam KP. Longitudinal assessment of an ELISPOT test for *Mycobacterium tuberculosis* infection. *PLoS Med.* 4(6):e192, 2007.

Pai M, Joshi R, Dogra S, Mendiratta DK, Narang P, Kalantri S, Reingold AL, Colford JM Jr, Riley LW, Menzies D. Serial testing of health care workers for tuberculosis using interferon-gamma assay. *Am J Respir Crit Care Med.* 174(3):349-55, 2006.

Menzies D. Interpretation of repeated tuberculin tests. Boosting, conversion, and reversion. *Am J Respir Crit Care Med.* 159(1):15-21, 1999.

Speaker #4

Implementation.....Masae Kawamura

Dewan PK, Grinsdale J, Kawamura LM. Low sensitivity of a whole-blood interferon-gamma release assay for detection of active tuberculosis. *Clin Infect Dis.* 1;44(1):69-73, 2007.

Dewan PK, Grinsdale J, Liska S, Wong E, Fallstad R, Kawamura LM. Feasibility, acceptability, and cost of tuberculosis testing by whole-blood interferon-gamma assay. *BMC Infect Dis.* 15;6:47, 2006.